



CARING FOR SOMEONE WITH COVID-19 AT HOME

Most people who get sick with COVID-19 have only mild illness, and should recover at home.* Care at home can help stop the spread of COVID-19, and help protect people who are at risk for getting seriously ill from COVID-19.

If you are caring for someone at home, monitor for emergency signs, prevent the spread of germs, treat symptoms, and carefully consider when to end home isolation.

*Note: Older adults and people of any age with certain serious underlying medical conditions like lung disease, heart disease, kidney disease or diabetes are at higher risk for developing more serious complications from COVID-19 illness, and should contact their health care provider as soon as symptoms start.

Monitor the person for worsening symptoms. Know the emergency warning signs.



- Have their healthcare provider's contact information on hand.
- If they are getting sicker, call their healthcare provider. For medical emergencies, call 911 and notify the dispatch personnel that they have, or are suspected to have, COVID-19.

People who develop emergency warning signs for COVID-19 should get medical attention immediately. Emergency warning signs include:*

- Difficulty breathing or shortness of breath
- Persistent pain or pressure in the chest
- New confusion or inability to arouse
- Bluish lips or face

*This list is not comprehensive. Please consult your medical provider for any other symptoms that are severe or concerning.

Provide symptom treatment.



- Make sure the sick person drinks a lot of fluids to stay hydrated and rests at home.
- Over-the-counter medicines may help with symptoms.
- For most people, symptoms last a few days and get better after a week.

CARING FOR SOMEONE WITH COVID-19 AT HOME



Prevent the spread of germs when caring for someone who is sick.



- Have the person stay in one room, away from other people, including yourself, as much as possible.
 - ◊ If possible, have them use a separate bathroom.
 - ◊ Avoid sharing personal household items, like dishes, towels and bedding.
 - ◊ If facemasks are available, have them wear a facemask when they are around people, including you.



- ◊ If the sick person can't wear a facemask, you should wear one while in the same room with them, if facemasks are available.
 - ◊ If the sick person needs to be around others (within the home, in a vehicle or at the doctor's office), they should wear a facemask.



- Wash your hands often with soap and water for at least 20 seconds, especially after interacting with the sick person. If soap and water are not readily available, use a hand sanitizer that contains at least 60% alcohol. Cover all surfaces of your hands and rub them together until they feel dry.



- Avoid touching your eyes, nose and mouth.
- Every day, clean all surfaces that are touched often, like counters, tabletops and doorknobs



- Use household cleaning sprays or wipes according to the label instructions.
- Wash laundry thoroughly.
 - ◊ If laundry is soiled, wear disposable gloves and keep the soiled items away from your body while laundering. Wash your hands immediately after removing gloves.
- Avoid having any unnecessary visitors.
- For any additional questions about their care, contact their health care provider or state or local health department.

When to end home isolation (staying home).



People **with COVID-19 who have stayed home (are home isolated)** can stop home isolation under the following conditions:

- **If they will not have a test** to determine if they are still contagious, they can leave home after these three things have happened:
 - ◊ They have had no fever for at least 72 hours (three full days of no fever without the use of medicine that reduces fevers), **AND**
 - ◊ Other symptoms have improved (for example, their cough or shortness of breath have improved), **AND**
 - ◊ At least 7 days have passed since their symptoms first appeared.
- **If they will be tested** to determine if they are still contagious, they can leave home after these three things have happened:
 - ◊ They no longer have a fever (without the use of medicine that reduces fevers), **AND**
 - ◊ Other symptoms have improved (for example, their cough or shortness of breath have improved), **AND**
 - ◊ They received two negative tests in a row, 24 hours apart. Their health care will follow CDC guidelines.