<u>Immunization Requirements for Preschool – 12th Grade</u>

3 Years	4 Years	5 Years	11-15 Years	16 Years and older
DTaP – 4 doses	*DTaP – 4 or 5 doses	*DTaP – 4 or 5 doses	*DTaP – 4 or 5 doses	*DTaP – 4 or 5 doses
Polio (IPV) – 3 doses	*Polio (IPV) – 3 or 4 doses	*Polio (IPV) – 3 or 4 doses	*Polio (IPV) – 3 or 4 doses	*Polio (IPV) – 3 or 4 doses
MMR – 1 dose	MMR – 2 doses	MMR – 2 doses	MMR – 2 doses	MMR – 2 doses
*Hib – 3 or 4 doses	*Hib – 3 or 4 doses	*Hib – no requirement	Tdap - 1 dose	Tdap – 1 dose
*Hep A - 2 doses effective for the school year beginning on or after July 1, 2018	*Hep A - 2 doses effective for the school year beginning on or after July 1, 2018	*Hep A - 2 doses effective for the school year beginning on or after July 1, 2018	*Hep A -2 doses effective for the school year beginning on or after July 1, 2018	*Hep A 2 doses effective for the school year beginning on or after July 1, 2018
Hep B – 3 doses	Hep B – 3 doses	Hep B – 3 doses	Hep B – 3 doses	Hep B – 3 doses
*Varicella – 1 dose	*Varicella – 2 doses	*Varicella – 2 doses	*Varicella – 2 doses	*Varicella – 2 doses
*Pneumococcal (PCV) – 4 doses	*Pneumococcal (PCV) – 4 doses	*Pneumococcal (PCV) – no requirement	Meningococcal ACWY (MCV4) – 1 dose (at 11 years of age)	*Meningococcal ACWY (MCV4) – 2 doses for 16 yrs and older effective for the school year beginning on or after July 1, 2018

^{*}DTaP - Fifth dose not needed if dose #4 given after 4th B-day and 6 months from previous dose

^{*} Hepatitis A – 2 doses spaced 6 months apart effective for the school year beginning on or after July 1, 2018





^{*} Polio - Fourth dose not needed if dose #3 given after 4th B-day and 6 months from previous dose

^{*} Hib - last dose must be on or after 12 months of age-not required at age 5 years and older; number of total doses (3 or 4) depends on brand used

^{*} PCV - last dose must be on or after 12 months of age-not required at age 5 years and older

^{*} Varicella - no doses needed if history of Chickenpox or Zoster disease is verified by a medical provider

^{*} Meningococcal ACYW – if initial dose given at 16 yrs or older, will only need one dose (previously known as MCV4)